



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Moreover the remaining factors will give you a *positive* product.
For, if one of the factors be

$$[(\xi_1 + i\eta_1) - (\xi_2 + i\eta)]^2 = [(\xi_1 - \xi_2) + i(\eta_1 - \eta_2)]^2$$

there will be a corresponding factor

$$[(\xi_1 - i\eta_1) - (\xi_2 - i\eta_2)]^2 = [(\xi_1 - \xi_2) - i(\eta_1 - \eta_2)]^2.$$

When multiplied these give

$$[(\xi_1 - \xi_2)^2 + (\eta_1 - \eta_2)^2]^2,$$

which is clearly positive. So all the factors containing complex roots may be taken in pairs which give a positive product, and, of course, the factors containing real roots only are positive. Hence corresponding to every pair of complex roots there will be *one and only one* negative factor. Therefore, if the number of pairs be even the discriminant will be positive; if odd, negative.

NOTE.—Mr. Harry S. Vandiver should have been credited as a joint author of the solution of problem 102.

GEOMETRY.

129. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Show that at no point of an ellipse will the circle of curvature pass through the center, if the eccentricity be less than $\frac{1}{2}\sqrt{2}$.

I. Solution by F. ANDEREGG, A.M., Professor of Mathematics, Oberlin College, Oberlin, O.; W.H. CARTER, A. M., Professor of Mathematics, Centenary College, Jackson, La.

Since for the ellipse the radius of curvature is

$$\rho = \frac{(a^4 y^2 + b^4 x^2)^{\frac{3}{2}}}{a^4 b^4};$$

and the center of curvature is the point

$$\left(\frac{(a^2 - b^2)x^3}{a^4}, -\frac{(a^2 - b^2)y^3}{b^4} \right),$$

the equation of the circle of curvature is

$$\left(x' - \frac{(a^2 - b^2)x^3}{a^4} \right)^2 + \left(y' + \frac{(a^2 - b^2)y^3}{b^4} \right)^2 = \frac{(a^4 y^2 + b^4 x^2)^3}{a^8 b^8}.$$

This circle passes through the origin

$$(a^2 - b^2)^2 (b^8 x^6 + a^8 y^6) = (a^4 y^2 + b^4 x^2)^3.$$